

# Imaging with Coherent X-rays: From Technique Development to Single Particles

*Sébastien Boutet<sup>a</sup>*

*<sup>a</sup>Linac Coherent Light Source, SLAC National Accelerator Laboratory*

*Author Email: sboutet@slac.stanford.edu*

The technique of Coherent X-ray Diffractive Imaging (CXDI) [1] utilizes algorithmic phase retrieval to generate an image of a sample using measured diffracted intensities. The achievable resolution is limited in principle only by the wavelength of the incident x-rays and the size of the detector used. It promises the ability to obtain a high-resolution image of any object, without the need for crystals. It will soon be two decades since the original demonstration of the use of coherent x-rays to image non-periodic objects [2]. Since this initial demonstration, a lot has changed in the world of x-ray science and not surprisingly, a lot of developments have taken place in the pursuit of high resolution coherent x-ray diffractive imaging.

Over the years, intense effort in algorithm development has made CXDI a fairly mature method that can now be employed in multiple fields such as biology [3] and condensed matter systems [4]. Scanning techniques have been developed allowing extended objects to be imaged. On overview of the CXDI technique and recent applications will be presented.

New x-ray sources are transforming x-ray science by enabling previously unachievable measurements to be performed. The new storage ring machines coming online will provide much increased coherent flux potentially allowing measurements that were previously signal-limited. Fourth generation light sources such as free electron lasers offer new and unique possibilities of limiting the radiation damage during the measurement [5] as well as unprecedented time resolution. The use of the CXDI technique with new x-rays sources will be presented, including time-resolved studies in materials [6] and developments towards using CXDI to image single particles, potentially single biological molecules [7].

## References

- [1] Miao, J.W., et al., Beyond crystallography: Diffractive imaging using coherent x-ray light sources. *Science*, 2015. **348**(6234): p. 530-535.
- [2] Miao, J.W., et al., Extending the methodology of X-ray crystallography to allow imaging of micrometre-sized non-crystalline specimens. *Nature*, 1999. **400**(6742): p. 342-344.
- [3] Nelson, J., et al., High-resolution x-ray diffraction microscopy of specifically labeled yeast cells. *Proceedings of the National Academy of Sciences of the United States of America*, 2010. **107**(16): p. 7235-7239.
- [4] Cha, W., et al., Core-shell strain structure of zeolite microcrystals. *Nature Materials*, 2013. **12**(8): p. 729-734.
- [5] Neutze, R., et al., Potential for biomolecular imaging with femtosecond X-ray pulses. *Nature*, 2000. **406**(6797): p. 752-757.
- [6] Clark, J.N., et al., Ultrafast Three-Dimensional Imaging of Lattice Dynamics in Individual Gold Nanocrystals. *Science*, 2013. **341**(6141): p. 56-59.
- [7] Hantke, M.F., et al., High-throughput imaging of heterogeneous cell organelles with an X-ray laser. *Nature Photonics*, 2014. **8**(12): p. 943-949.